

A CHECKLIST OF THE NATIVE ANNUAL FLORA OF CONTINENTAL CHILE

UN CATALOGO DE LA FLORA ANUAL NATIVA DE CHILE CONTINENTAL

Mary T. Kalin Arroyo*, Clodomiro Marticorena ** and Melica Muñoz***

ABSTRACT

A checklist of the native annual species of continental Chile is provided. Observations on 4408 native species (over 99% of the total native vascular flora of continental Chile) indicate the presence of the annual habit in 692 species (15.7% of the flora). Annuals occurs in 196 genera (24.7%) and 53 families (31.2%). There are proportionately more annuals among dicotyledons (19.3%) than among monocotyledons (4.3%), a trend also seen to be statistically significant at the generic level. 358 (51.7%) annual species are fairly certainly endemic to continental Chile, while an additional 18 species (2.6%) are characterized by endemic Chilean infraspecific taxa. A further 20 species are probably endemic to continental Chile as are varieties of 3 species. Seven of Chile's endemic genera (10.4%) are exclusively annual (*Cyphocarpus* (Campanulaceae); *Microphytes* (Caryophyllaceae); *Moscharia* (Asteraceae); *Agallis* (Cruciferae); *Scyphanthus* (Loasaceae); *Homalocarpus* (Umbelliferae); *Araeoandra* (Vivianaceae)). The frequency of the annual habit in Chile is briefly compared with that of other arid and semi-arid regions.

KEYWORDS: Annual habit; flora of continental Chile; endemic species and genera; mediterranean-type climate.

RESUMEN

Se presenta un catálogo de las especies anuales de la flora nativa de Chile continental. Observaciones en 4408 especies (más del 99% de la flora vascular nativa total de Chile continental) indica la presencia del hábito anual en 692 especies (15,7% de la flora). Las anuales pertenecen a 196 géneros (24,7%) y 53 familias (31,2%). La frecuencia del hábito anual es significativamente mayor en las dicotiledóneas (19,3%) que en las monocotiledóneas (4,3%), una tendencia que también es estadísticamente significativa al nivel de género. 358 (51,7%) de las especies anuales son endémicas a Chile continental, en tanto que 18 (2,6%) especies presentan variedades chilenas endémicas. 20 especies adicionales probablemente son endémicas tal como lo son variedades de 3 especies adicionales. Siete de los géneros endémicos de Chile continental (10,4% del total) son exclusivamente anuales (*Cyphocarpus* (Campanulaceae); *Microphytes* (Caryophyllaceae); *Moscharia* (Asteraceae); *Agallis* (Cruciferae); *Scyphanthus* (Loasaceae); *Homalocarpus* (Umbelliferae); *Araeoandra* (Vivianaceae)). Se compara la frecuencia del hábito anual en la flora nativa de Chile continental con la de otras áreas áridas y semi-áridas.

INTRODUCTION

* Departamento de Biología, Facultad de Ciencias, Universidad de Chile, Casilla 653, Santiago, Chile.

** Departamento de Botánica, Facultad de Ciencias Biológicas y de Recursos Naturales, Universidad de Concepción, Casilla 2407, Chile.

*** Sección Botánica, Museo Nacional de Historia Natural, Santiago, Chile.

A significant proportion of continental Chile is characterized by arid to semi-arid climates with strongly seasonal or highly irregular rainfall (di Castri & Hajek, 1976). Desert to semi-desert climates extend along the coast and far inland to above 2000 m from the far north (17°S) to

around 26°S (Arroyo et al., 1988). In central Chile from 32° - 38°S and in interior valleys as far south as 40°S the climate is typically mediterranean (di Castri, 1988). A semi-mediterranean-type climate appears in the extreme eastern border of the Chilean Patagonia (di Castri & Hajek, 1976).

Many temperate semi-arid and arid regions are known to support taxonomically diverse and speciose native annual assemblages. The State of California, with a predominantly mediterranean-type climate, supports 28.6% native annuals (Raven & Axelrod, 1978) while the Sonoran desert is characterized by 21.4% (Shreve & Wiggins, 1964). The flora of Israel, with only 1.7% aliens across all life-forms, supports more than 50% annuals (Eig, 1931, 1932; Danin & Orshan, 1990). A notable exception to this rule, is the mediterranean-type climate Cape region of South Africa with only 6.4% native annuals (Goldblatt, 1978). Vegetation studies also suggest a relatively low percentage of annuals for mediterranean-type climate areas in southern Australia (Adamson, 1927). Interestingly, both mediterranean-type climate areas with lower percentages of annuals are found in the southern hemisphere.

Chile, mainly as a result of a lack of a comprehensive floristic treatment for the country, is one of the last mediterranean-type climate regions to be surveyed for the annual habit. Some data has been forthcoming from vegetation plots, small scale floristic studies and other ecological studies (e.g. Schlegel, 1966; Mooney et al., 1977; Arroyo & Squeo, 1990; Arroyo & Uslar, 1990). Additional information for a few areas may be gleaned from regional floras (e.g. Moore, 1982; Arroyo et al., 1984; Arroyo et al., 1988; Arroyo et al., 1989). However the present state of knowledge of the frequency of the annual habit is insufficient for characterizing the Chilean flora in general.

Here we provide a complete checklist of the native annual species of continental Chile. The information has been gathered in relation to a more comprehensive study of similarities and differences between the vascular floras of Chile and California, to be presented elsewhere. Given the lack of a modern flora, publication of a checklist of the native annuals of Chile seems timely. Such information should be useful to professional

plant ecologists who are presently forced to work with limited botanical literature. Secondly, agro-nomists interested in native forage crops should benefit from the availability of such a checklist. The checklist has been compiled to a large degree from literature sources. In that some authors give more attention to life-form than others, further observations by field biologists using it constitute the best avenue for its perfection.

INFORMATION SOURCES

To assess the incidence of the annual habit in continental Chile a data base of all native vascular plant species occurring there was developed. By continental Chile we mean *mainland Chile excluding the Juan Fernández Islands, the San Ambrosio and San Félix Islands and Easter Island*, all under Chilean jurisdiction. The data base relied on the checklist of the Chilean flora published by Marticorena & Quezada (1985), as baseline information, modified as follows. Species in the Marticorena & Quezada (1985) checklist restricted to the forementioned island territories of Chile and all introduced species in the list were excluded. The reduced list was further emended for new species, numerous synonymy changes published after 1985 and corrections of the alien status of some species. Infraspecific taxa are not considered. The final data base contained 4443 native vascular plants species distributed in 793 genera and 170 families.

For assessing the annual habit the relevant life-form information for many species was obtained directly from recent generic revisions. For others we found it necessary to rely on the older Reiche (1896-1911) flora, local floras for Argentina and Chile (e.g. Moore, 1982; Correa, 1971) and original species descriptions. Such information was complemented with our own field knowledge of life-forms and consultation of herbarium material.

Fairly reliable information was found for a total of 4408 (>99% of the 4443) names in the continental checklist. For the present purposes, 20 species, given as "probably annual" are given the benefit of the doubt. Species varying in habit from an annual to biennial or perennial herb are designated as facultatively annual. Annual spe-

cies and genera with annuals endemic to continental Chile are indicated.

CHECKLIST OF THE NATIVE ANNUALS OF CONTINENTAL CHILE

(fa) following the specific epithet indicates a *facultative annual*. *Probable annuals* are followed by (a?). All other species are considered *obligate annuals*. Annual species *endemic* to continental Chile are indicated by (e). Annual species which contain an *endemic variety* in Chile are indicated by (ev). Genera endemic to Chile with annuals are designated by (E). The questioned endemics (?) are unsure endemics. (nv) following a specific epithet indicates a species occurring in Chile represented by a native variety in addition to an alien variety.

PTERIDOPHYTA

SALVINIACEAE

Azolla

filiculoides Lam.

ANGIOSPERMAE: DICOTYLEDONEAE

AIZOACEAE

Tetragonia

copiapina Phil. (e)
espinosae Muñoz (e)
macrocarpa Phil. (ev)
pedunculata Phil.
tenella Johnst. (e)
trigona Phil. (fa)(e)

AMARANTHACEAE

Amaranthus

asplundii Thell.
looseri Suess. (e)

Gomphrena

umbellata Remy

ASTERACEAE

Agoseris

chilensis (Less.) Greene (e)
coronopifolia (D'Urv.) Chamb. ex D.M. Moore (fa)

Amblyopappus

pusillus H. et A.

Bidens

andicola Kunth. (fa)

Blennosperma

chilense Less. (e)

Chaetanthera

aymarae Martic. et Quez. (e)
ciliata R. et P. (e)
euphrasioides (DC.) Meigen
flabellata D. Don (e)
flabellifolia Cabr. (e)
glabrata (DC.) Meigen (e)
gnaphalioides (Remy) Johnst.
incana Poepp. ex Less. (e)
leptocephala Cabr. (e)
limbata (D. Don) Less. (e)
linearis Poepp. ex Less. (e)
microphylla (Cass.) H. et A. (ev)
minuta (Phil.) Cabr.
moenchioides Less. (e)
planiseta Cabr. (e)
pusilla (D. Don) H. et A.
splendens (Remy) B.L. Rob.
tenella Less. (e)

Conyza

artemisiifolia Meyen et Walp.
bustillosiana Remy (e)
copiapina Phil. (a?)(e)
floribunda H.B.K. (fa)
glabrata Phil. (fa)(e)
hirtella (DC.) Martic. (e)*
lechleri (Sch. Bip.) Cabr.
minutiflora Phil. (e)

* *Conyza hirtella* (De Candolle) Marticorena, nov. comb.
Basionimo: *Erigeron hirtellus* De Candolle, Prodr. 5: 290.
1836.

Cotula

mexicana DC.

Doniophyton

anomalum (D. Don) Kurtz
patagonicum (Phil.) Hieron.

Eclipta

prostrata (L.) L. (fa)

Erechtites

leptantha (Phil.) Cabr. (e)

Facelis

plumosa (Wedd.) Sch. Bip.
retusa (Lam.) Sch. Bip. (ev)

Flaveria

bidentis (L.) O.K.

Galinsoga

parviflora Cav.
quadriradiata R. et P.

Gamochaeta

monticola (Phil. ex Reiche) Cabr.
simplicicaulis (Willd. ex Sprengel) Cabr.
spachelata (H.B.K.) Cabr.
stachydifolia (Lam.) Cabr.

Gnaphalium

aldunateoides Remy
cheiranthifolium Lam. (fa)
cymatoides Kunze ex DC. (ev)
diminutivum Phil. (e?)
heterotrichum Phil. (e)
moelleri Phil.
perpusillum Phil. (e?)
phaeolepis Phil. (e)
pratense Phil.
ramosum Phil. (e)

Helenium

aromaticum (Hook.) Bailey (fa)(e)
atacamense Cabr. (fa)(e)
ovallense Bierner (fa)(e)
urmenetae (Phil.) Cabr. (fa)(e)
vallenariense (Phil.) Bierner (fa)(e)

Heterosperma

nanum (Nutt.) Sherff
ovatifolium Cav.

Hieracium

antarcticum D'Urv.

Lasthenia

kunthii (Less.) H. et A. (e)

Leucheria

cerberoana Remy (e)
cumingii H. et A. (e)
glabriuscula (Phil.) Reiche (e)
glandulosa D. Don (e)
menana Remy (e)
oligocephala Remy (e)
seneciooides H. et A. (e)
tenuis Less. (e)

Madia

chilensis (Nutt.) Reiche (e)
sativa Mol.

Malacothrix

chevelandii A. Gray
coultieri A. Gray

Micropsis

nana D.C.

Microseris

pygmaea D. Don

Moscharia (E)

pinnatifida R. et P. (e)
solbrigii Crisci (e)

Polyachyrus

annuus Johnst.

Psilocarphus

brevissimus Nutt.

Schkuhria

multiflora H. et A.
pinnata (Lam.) O.K.

Senecio

troncosii Phil. (e)

Sigesbeckia

jurullensis H.B.K.

Soliva

pterosperma (A.L. Juss.) Less.
sessilis R. et P. (e)
stolonifera (Brot.) Loud.
valdiviana Phil.

Tagetes

biflora Cabr.
minuta L.
multiflora H.B.K.

Triptilion

achilleae DC. (fa)
berteroii Phil. (a?)(e)
capillatum (D. Don) DC. (e)
cordifolium Lag. ex Lindl. (e)
diffusum (D. Don) DC. (a?)(e)
digitatum Phil. (e)
dusenii O. Hoffm. (e)
euphrasioides Bert. ex DC. (e)
gibbosum Remy (e)

Verbesina

encelioides (Cav.) B. et H. ex A. Gray (fa)

Villanova

oppositifolia Lag.

Xanthium

argenteum Widder (e)

BORAGINACEAE

Amsinckia

calycina (Moris) Chater
tessellata A. Gray

Cryptantha

alfalfalis (Phil.) Johnst. (e)
aspera (Phil.) Grau (e)
calycina (Phil.) Reiche (e)
calycotricha Johnst. (e)
chaetocalyx (Phil.) Johnst. (e)
chispae Grau (e)
clandestina (Trev.) Johnst. (e)
cynoglossoides (Phil.) Johnst.
dichita (phil.) Johnst. (e)

difussa (Phil.) Johnst.

dimorpha (Phil.) Greene (e)

diplotricha (Phil.) Reiche

dichophylla (Phil.) Reiche (e)

filaginea (Phil.) Reiche

filiformis (Phil.) Reiche (e)

gayi Johnst. (fa)(e)

globulifera (Clos) Reiche

glomerata Lehm. (e)

haplostachya (Phil.) Johnst. (e)

hispida (Phil.) Reiche (e)

involucrata (Phil.) Reiche (fa)(e)

kingii (Phil.) Reiche (e)

marticorenae Grau (e)

parviflora (Phil.) Reiche

phaceloides (Clos) Reiche (e)

romanii Johnst. (e)

taltalensis Johnst. (e)

volckmannii (Phil.) Johnst. (e)

werdermanniana Johnst. (e)

Heliotropium

geissei F. Phil. (e)

microstachyum R. et P.

paronychioides A. DC.

Myosotis

antarctica Hook. f. (fa)

Pectocarya

anomala Johnst.

dimorpha (Johnst.) Johnst.

linearis (R. et P.) DC.

pusilla (A. DC) A. Gray

Plagiobothrys

armeriifolius (Phil.) Johnst. (e)

calandrinooides (Phil.) Johnst.

collinus (Phil.) Johnst. (e)

corymbosus (R. et P.) Johnst. (e)

fulvus (H. et A.) Johnst. (e)

gracilis (R. et P.) Johnst. (e?)

myosotoides (Lehm.) Brand

oppositifolius (Phil.) Johnst. (e)

polycaulis (Phil.) Johnst. (fa)(e)

pratensis (Phil.) Johnst. (e)

procumbens (Colla) A. Gray

pulchellus (Phil.) Johnst. (e)

uliginosus (Phil.) Johnst. (e)

verrucosus (Phil.) Johnst.

CALLITRICHACEAE

Callitricha

antarctica Engelm, ex Hegelm.
hermafroditica L.
lechleri (Hegelm.) Fassett
palustris L.
terrestris Rafin.

CALYCERACEAE

Boopis

gracilis Phil.
pusilla Phil. (e)

Calycera

eryngioides Remy
integrifolia (Phil.) Reiche (e)
leucanthema (Poepp.) Reiche (e?)
sessiliflora Phil. (e)

Moschopsis

monocephala (Phil.) Reiche (fa)

CAMPANULACEAE

Cyphocarpus (E)

innocuous Sandw. (e)
psammophilus Ricardi (e)
rigescens Miers (e)

Downingia

pusilla (Poepp. ex Cham.) Torr.

Legenere

valdiviana (Phil.) Wimmer (fa)

Lobelia

alata Labill.

Triodianis

biflora (R. et P.) Greene

CAPPARACEAE

Cleome

chilensis DC. (ev?)

CARYOPHYLLACEAE

Arenaria

oligosperma Naud. (a?)(e?)

Drymaria

cordata (L.) Willd. ex Roem. et Schult.
engleriana (Muschl.) Baehni et MacBr.
paposana Phil. (ev)

Microphyes (E)

litoralis Phil. (e)
minimus (Bertero ex Colla) (Briq.) (e)
robustus Ricardi (e)

Minuartia

acutiflora (Fenzl) Mattf.

Sagina

apetala Ard. (ev?)
chilensis Naud.

Spergularia

cremnophila Johnst. (fa)(e)
denticulata Phil. (e)
platensis (Cambess) Fenzl
stenocarpa (Phil.) Johnst. (fa)

Stellaria

abortiva Naud. (e?)

CHENOPODIACEAE

Atriplex

chilensis Colla (e)
myriophylla Phil.
oreophila Phil.
peruviana Moq.
philippii R. E. Fries (e)

Chenopodium

ambrosioides L. (fa)
antarcticum (Hook. f.) Hook. f.
carnosulum Moq.
frigidum Phil.
macrospermum Hook. f.
papulosum Moq.
petiolare H.B.K.
philippianum Aellen
quinoa Willd.

Nitrophila

atacamensis (Phil.) Hieron. ex Ulbr. (fa)

Suaeda

patagonica Speg.

CRASSULACEAE

Crassula

closiana (Gay) Reiche (e)
connata (R. et P.) Berger
decumbens Thunb.
moschata G. Forster
ovallei (Phil.) Reiche (e)
peduncularis (J.E. Sm.) Meigen
solieri (Gay) Meigen
tillaea Lest.-Garl.

CRUCIFERAE

Agallis (E)

lanata (Barn.) Gilg et Muschl. ex O.E. Schultz (e)

Cardamine

chilensis (e?) DC.
nivalis Gill. ex Hook. et Arn. (e)
solisii Phil. (e?).

Coronopus

leptocarpus Boelcke (e)

Cremolobus

chilensis (Lag. ex DC.) DC.

Descurainia

cumingiana (Fisch. et Mey.) Prantl (fa)(ev)
diversifolia O.E. Schulz (e)
glaucescens (Phil.) O.E. Schulz (fa)
nuttallii (Colla) O.E. Shulz (a?)(e)
pinnata (Walter) Britton
stricta (Phil.) Prantl ex Reiche (fa)(ev)

Diplotaxis

chilensis Barn. (e)

Draba

australis R. Br.

Lepidium

ales Macbr.
angustissimum Phil. (e)
auriculatum Regel et Koern. (fa)(e)
bipinnatifidum A.N. Desv. (fa)
brevicaule Barn. (e)
curicoanum Phil. (e)
johnstonii C. Hitch. (e)
myrianthum Phil.
nidum Nutt. ex Torr. et Gray (e)
pseudo-didymus Thell. ex Druce (fa)
pubescens A.N. Desv.
rahmeri Phil. (fa)
raimondii O.E. Schulz
strictum (S. Wats.) Rattan.
subvaginatum Steud. ex Thell. (fa)

Menonvillea

chilensis (Turcz.) Jacks. (e)
filifolia Fischer et C. Meyer (fa)(e)
gayi Phil.
linearis DC. (fa)(e)
litoralis (Barn.) Rollins (e)
minima Rollins (e)
orbiculata Phil. (fa)(e)
pinnatifida Barn. (fa)(e)

Rorippa

austroamericana Mart.-Lab. (fa)
philippiana (Speg.) Maclosk. (fa)

Schizopetalon

arcuatum Al-Shehbaz (e)
bipinnatifidum Phil. (e)
biseriatum Phil. (e)
brachycarpum Al-Shehbaz (e)
corymbosum Al-Shehbaz (e)
dentatum (Barn.) Gilg et Muschl. (e)
maritimum Barn. (e)
rupestre (Barn.) Reiche
tenuifolium Phil. (e)
walkeri Hook. (e)

Thlaspi

alpestre (ev?)

ELATINACEAE

Elatine

triandra Schkuhr (fa)

EUPHORBIACEAE

Euphorbia

- germainii Phil. (e)
- meyeniana Klotzsch (fa)
- minuta Phil.
- pygmaea Phil. (e)
- tarapacana Phil. (e)
- verna Phil. (e)

GENTIANACEAE

Centaurium

- cachanalahuensis (Mol.) B.L. Rob.

Cicendia

- quadrangularis (Lam.) Griseb.

Gentiana

- lactea Phil. (e)
- prostrata Haenke (fa)

Gentianella

- magellanica (Gaud.) Fabris ex D.M. Moore (fa)

GERANIACEAE

Geranium

- intermedium Colla, ex Savi (e)

HYDROPHYLACEAE

Nama

- dichotomum (R. et P.) Choisy
- undulatum H.B.K. (ev).

Phacelia

- brachyantha Benth.
- cummingii (Benth.) A. Gray
- nana Wedd.
- pinnatifida Griseb. ex Wedd. (fa)
- setigera Phil.

LABIATAE

Salvia

- paposana Phil.

Stachys

- eremicola Epling (e)
- gilliesii Benth. (fa)
- truncata Kunze ex Benth. (e)

LOASACEAE

Caiophora

- contorta (Desr.) Urban et Gilg
- dissecta (Hook.) Urban et Gilg (fa)(e)
- tomentosula Urban et Gilg (fa)(e)

Loasa

- aphanantha Urban et Gilg
- artemisiifolia (Poepp.) Urban et Gilg (fa)(e)
- bertrandii Phil. (e)
- caespitosa Phil. (e)
- chilensis (Gay) Urban et Gilg (e)
- floribunda H. et A. (e)
- gayana Urban et Gilg (e)
- intricata Gay (e)
- lateritia Gill. ex Arn.
- longiseta Phil. (e)
- malesherbioides Phil. (e)
- martinii Phil. (e)
- micrantha Poepp. (e)
- multifida Gay (a?)(e)
- pallida Gill. ex Arn. (e)
- tricolor Ker-Gawl. (ev)
- triloba Domb, ex A.L. Juss. (e)
- urens Jacq.
- urmenetae Phil. (e)
- volubilis Domb. ex A.L. Juss. (fa)(e)

Mentzelia

- bartonoides (K. Presl) Urban et Gilg
- pinnatifida (Phil.) Urban et Gilg
- solieri (Gay) Urban et Gilg

Scyphanthus (E)

- elegans D. Don (fa)(e)
- stenocarpus (Poepp.) Urban et Gilg (fa)(e)

LYTHRACEAE

Pleurophora

- polyandra H. et A. (e)
- pusilla H. et A. (e)

MALESHERBIACEAE

Malesherbia

- gabrielae Ricardi (e)
humilis Poepp. (e)
multiflora Ricardi (e)
taltalina Ricardi (e)

MALVACEAE

Cristaria

- adenophora Johnst. (e)
australis Phil. (e)
cyanea Phil ex E. Baker (e)
dissecta H. et A.
divaricata Phil. ex E. Baker
diversifolia Phil. (e)
elegans Gay (a?)(e)
eriantha H. et A. (fa)(e)
flexuosa Phil. (a?)(e)
formosula Johnst.
glandulosa Phil. (e)
heterophylla (Cav.) H. et A. (e)
hirsuta K. Presl (e)
humilis Phil. (e)
inconspicua F., Phil. ex Phil. (e)
integerrima Phil. (fa)(e)
intermedia Gay (fa)(e)
intonsa Johnst. (e)
molinae Gay (e)
ovallea Gay (e)
patens Phil. (e)
pinnatifida H. et A. (e)
ranunculifolia Phil. ex E. Baker (e)
rotundifolia Phil. (e)
saniculifolia Phil. ex E. Baker (e)
sundtii Phil. (e)
trifida Phil. (e)
univittata Hochr. (e)

Lecanophora

- heterophylla (Cav.) Krap. (e)

Nototriche

- diminutiva (Phil.) Johnst. (e)
nana A.W. Hill
pusilla A.W. Hill
pygmaea (Remy) A.W. Hill
sarmentosa A.W. Hill

Palaua

- inconspicua Johnst.
modesta (Phil.) Reiche (e)

Tarasa

- antofagastana (Phil.) Krap.
congestiflora (Johnst.) Krap.
tarapacana (Phil.) Krap.
tenella (Cav.) Krap.
umbellata Krap. (e)

Urocarpidium

- chilense (Braun et Bouché) Krap.
peruvianum (L.) Krap.

MOLLUGINACEAE

Glinus

- radiatus (Ruiz et Pavón) Rohrb.

NOLANACEAE

Nolana

- aplocaryoides (Gaud.) Johnst. (e)
baccata (Lindl.) Dunal (e)
gracillima Johnst.
intonsa I.M. Johnst. (fa)(e)
parviflora (Phil.) Phil. (e)
pterocarpa Phil. ex Wettst. (e)
rhombifolia Martic. et Quez. (e)

ONAGRACEAE

Camissonia

- dentata (Cav.) Reiche (ev)

Clarkia

- tenella (Cav.) Lewis et Lewis (ev)

Gayophytum

- humile A.H.L. Juss.
micranthum H. et A.

Oenothera

- affinis Cambess.
arequipensis Munz et Johnst.
coquimbensis Gay (e)
grisea W. Dietr. (e)

- magellanica Phil. (fa)
nana Griseb. (fa)
odorata Jacq.
peruana W. Dietr. (fa)
picensis Phil.
ravenii W. Dietr. (fa)(ev)
rubida Rusby
sandiana Hassk.
villaricæ W. Dietr. (fa)

OXALIDACEAE

Oxalis

- aberrans Reiche (e)
clandestina Phil. (fa)(e)
compacta Gill. ex Hook. et Arn. (fa)(ev)
laxa H. et A.
leptocaulos Phil. (fa)(e)
ovalleana Phil. (e)
pubescens H.B.K.
rigida (Barn.) Lourt.
rosea Jacq. (e)
valdiviensis Barn. (fa)

PAPAVERACEAE

Argemone

- hunnemannii Otto et Dietr. (fa)(e)
rosea Hook. (e)
subfusiformis Ownbey

PAPILIONACEAE

Adesmia

- capitellata (Clos) Haum.
eremophila Phil. (fa)(e)
filifolia Clos (e)
micrantha Phil. (e)
multicuspis Clos (e)
muricata (Jacq.) DC.
parviflora Clos (e)
pusilla Phil. (e)
rahmeri Phil. (fa)
tenella H. et A. (e)

Astragalus

- berteroanus (Moris) Reiche (e)
coquimbensis (H. et A.) Reiche (e)

- dodtii Phil. (e)
paposanus Johnst. (e)
pissisii (phil.) Johnst. (e)
triflorus (DC.) A. Gray

Dalea

- moquehuana Macbr.

Lathyrus

- campestris Phil.
crassipes Gill. ex H. et A.

Lotus

- subpinnatus Lag.

Lupinus

- microcarpus Sims

Trifolium

- chilense H. et A. (e)
circundatum Kunze (e)
depauperatum A.N. Desv.
macraei H. et A.
microdon H. et A.
physanthum H. et A. (e)
triaristatum Bert. et Savi
vernun Phil. (e)

Vicia

- acerosa Clos
berteroana Phil. (a?)(e?)
ciliaris Phil. (a?)(e?)
graminea J.E. Sim.
inconspicua Phil. (e?)
leyboldii Phil. (fa)(e?)
magnifolia Clos (e?)
micrantha H. et A. (e?)
modesta Phil. (e?)
moorei Phil. (a?)(e?)
sessiliflora Clos (a?)(e?)
subserrata Phil. (e?)
truncata Phil. (a?)(e?)
vicina Clos (e?)

PLANTAGINACEAE

Plantago

- firma Kunze et Walp.
hispidula R. et P. (e)
limensis Pers.

litorea Phil. (e)
pulvinata Speg. (fa)
rancagua Steud. (e)

POTULACACEAE

POLEMONIACEAE

Collomia

biflora (R. et P.) Brand
cavanillesii H. et A. (e)

Gilia

crassifolia Benth.
glutinosa Phil.
laciniata R. et P.
valdiviensis Griseb.

Ipomopsis

gossypifera (Gill. ex Benth.) Grant

Linanthus

pusillus (Benth.) Greene (e)

Microsteris

gracilis (Dougl. ex Hook.) Greene

Navarretia

involuta R. et P.

Polemonium

micranthum Benth

POLYGALACEAE

Polygala

gayi A. Benn. (fa)(e)

Calandrinia

acuminata Phil. (e)
arenaria Cham. (e)
banduriae Phil.
berteroana Phil. (e)
cachinalensis Phil. (e)
cephalophora Johnst. (e)
chrysantha Johnst. (e)
ciliata (R. et P.) DC. (a?)
compressa Schrad. ex DC. (e)
coquimbensis Barn. (e)
cumingii H. et A. (e)
cymosa Phil. (e)
demissa Phil. (e)
densiflora Barn.
erythrocoma Phil. (e)
glaucopurpurea Reiche (e)
glomerata Phil. (e)
litoralis Phil. (e)
longiscapa Barn. (e)
modesta Phil.
oblongifolia Barn. (e)
parviflora Phil. (e)
polycarpoides Phil.
ramosissima H. et A. (e)
speciosa Lehm. (e)
spicata Phil. (e)
spicigera Phil. (e)
stricta Phil. (e)
thyrsoides Reiche (e)
trifida H. et A. (e)
villanuevae Phil. (a?)(e)

POLYGONACEAE

Chorizanthe

commisuralis Remy (e)

Lastariaea

chilensis Remy (e)

Oxytheca

dendroidea Nutt.

Monocosmia

monandra (R. et P.) Baillon

Montia

fontana L.

Philippia

amaranthoides (Phil.) O.K. (e)
celosioides (Phil.) O.K. (e)
fastigiata (Phil.) Pax ex K. Hoffm. (e)

PRIMULACEAE

Anagallis

minima (L.) Krause

Androsace
salasii Kurtz

Pelletiera
verna St.-Hil.

RANUNCULACEAE

Myosurus
apetalus Gay
patagonicus Speg.

Ranunculus
apiifolius Pers.
bonariensis Poir.
chilensis DC.
flagelliformis J.E. Sm.
pseudotrullifolius Skottsb.

ROSACEAE

Aphanes
berteroana Rothm. (e)
looseri (Rothm.) Rothm. (e)
neglecta (Rothm.) Rothm.

RUBIACEAE

Cruckshanksia
pumila Clos (e)
tripartita Phil. (fa)(e)

Galium
fuegianum Hook. f. (fa)
aparine L.**

SANTALACEAE

Quinchamalium
berteroanum Phil. (e)
bracteosum Phil. (e)
carnosum Phil. (fa)(e)
excrescens Phil. (e)
litorale Phil. (e)
parviflorum Phil. (e)

** The status of *Galium aparine* L. in Chile is complex. Dempster (1981) states "it is highly probable that South American plants are for the most part native, but that the species has also been introduced from Europe...".

SAXIFRAGACEAE

Lepuropetalon
spathulatum (Muhl.) Elliot

SCROPHULARIACEAE

Bartsia
chilensis Benth. (e)

Calceolaria
bipinnatifida Phil.
hollermayeri Kraenzl. (e)
scabiosifolia Sims

Euphrasia
antarctica Benth.
meiantha Clos (fa)
perpusilla Phil. (e)

Limosella
australis R. Br.

Lindernia
dubia (L.) Pennell (fa)

Mimulus
acaulis Phil. (e)
bridgesii (Benth.) Clos (e)
depressus Phil. (e)
glabratus H.B.K.
luteus L. (fa)(ev)
nanus Phil.

Orthocarpus
attenuatus A. Gray
lacinatus (H. et A.) Keck

SOLANACEAE

Cacabus
flavus Johnst.
integrifolius Phil. (e)

Nicandra
physalodes (L.) Gaertn.

Nicotiana
acuminata (Graham) Hook. (ev)

corymbosa Remy
linearis Phil.
longibracteata Phil.
miersii Remy (e)
noctiflora Hook. (fa)
pauciflora Remy (e)
petunioides (Griseb.) Millan
undulata R. et P.

Reyesia

parviflora (Phil.) Hunz.

Schizanthus

alpestris Poepp. ex Benth. (e)
candidus Lindl. (fa)(e)
grahamii Gill. ex Hook.
hookeri Gill. ex Graham (fa)(e)
integrifolius Phil. (e)
lacteus Phil. (e)
laetus Phil. (e)
litoralis Phil. (e)
parvulus Sudzuki (e)
pinnatus R. et P. (e)
porrigens Graham (e)
tricolor Grau et Gronb. (e)

Solanum

andinum Reiche (a?)(e)
euacanthum Phil.
furcatum Dunal ex Poir.
gaudichaudii Dunal (e)
heterantherum Witasek ex Reiche (e)
maritimum Meyen ex Nees (e)
nigrum L. (ev)
patagonicum Morton (fa)
pentlandii Dunal

UMBELLIFERAE

Apium

laciniatum (DC.) Urban (ev)
leptophyllum (Pers.) F. Muell. ex Benth.

Asteriscium

aemocarpon Clos (e)
closii (O.K.) Math. et Const. (e)

Bowlesia

incana R. et P.
macrophysa Zoell. (e)

paposana Johnst. (e)
sodiroana Wolff
uncinata Colla (e)

Daucus

montevidensis Link ex Sprengel (a?)

Domeykoa

oppositifolia Phil. (e)

Eryngium

anomalum H. et A. (e)
coquimbanum Phil. ex Urban (e)
depressum H. et A. (e)
macracanthum Phil. (e)
pulchellum Phil. (e)

***Homalocarpus* (E)**

bowlesioides H. et A. (e)
dichotomus (Poepp. ex DC.) Math. et Const. (e)
digitatus (Phil.) Math. et Const. (e)
dissectus Math. et Const. (e)
integerrimus (Turcz.) Math. et Const. (e)
nigripetalus (Clos) Math. et Const. (e)

URTICACEAE

Parietaria

debilis G. Forster

Urtica

berteroana Phil.
flabellata H.B.K.

VALERIANACEAE

Plectritis

samolifolia (DC.) Hoeck (e)

Valeriana

aequiloba Clos (e)
crispa R. et P. (e)
floribunda Phil. (e)
grandifolia Phil. (e)
oreocharis Phil. (e)
polemoniifolia Phil.
sphaerocarpa Phil. (e)
valdiviana Phil. (e)
virescens Clos

VERBENACEAE

Verbena

dissecta Willd. (a?)

VIOLACEAE

Viola

araucaniae Becker (e)

aurata Phil. (e)

auricula Leyb. (e)

bicolor Reiche (e)

brachypetala Gay (a?)(e)

calderensis Becker (e)

chamaedrys Leyb. (e)

chrysantha Phil.

domeykoana Gay

frigida Phil. (ev)

glechomoides Leyb. (e)

godoyae Phil. (e)

johnstonii Becker (e)

litoralis Phil. (e)

llulliaillacoensis Becker (e)

minutiflora Phil. (e)

nubigena Leyb. (e)

ovalleana Phil. (e)

polypoda Turcz. (e)

pulchella Leyb. ex Reiche (e)

pulvinata Reiche (e)

pusilla Poepp. (e)

rhombifolia Leyb. (e)

taltalensis Becker (e)

vallenarensis Becker (e)

werdermannii Becker (e)

VIVIANACEAE

Araeoandra (E)

tenuicaulis (Barn.) Lefor (e)

ANGIOSPERMAE:

MONOCOTYLEDONEAE

AMARYLLIDACEAE

Alstroemeria

graminea Phil. (e)

CYPERACEAE

Cyperus

rivularis Kunth

volckmannii Phil. (e)

Scirpus

cernuus Vahl (nv)

GRAMINEAE

Agrostis

gelida Trin. (e)

oligoclada Phil. (e)

serranoi Phil. (e)

Alopecurus

heleochloides Hackel (e)

Bouteloua

simplex Lag.

Bromidium

anomalum (Trin.) Doell

trisetoides (Steud.) Rugolo (e)

Bromus

berterianus Colla

gunckelii Matthei (e)

Chaetotropis

chilensis Kunth

Deschampsia

airiformis (Steud.) Benth.

berteroana (Kunth) Trin. (e)

danthonioides (Trin.) Munro ex Benth. (e)

looseriana Parodi (e)

monandra Parodi (e)

Diplachne

uninervia (J. Presl) Parodi

Eragrostis

mexicana (Hornem.) Link

virescens J. Presl

Hordeum

pusillum Nutt.

Koeleria

grisebachii Domin

JUNCACEAE

Muhlenbergia

peruviana (P. Beauv.) Steud.

Juncus

bufonius L.

capitatus Weigel

Munroa

andina Phil.

decumbens Phil.

LILAEACEAE

Phalaris

amethystina Trin.

angusta Nees ex Trin.

Lilaea

scilloides (Poir.) Haum.

Poa

pumila Phil. (a?)

CONCLUDING REMARKS

Polypogon

linearis Trin. (e)

692 (15.7%) of the native vascular plant species of continental Chile are annual or facultatively annual (Table 1). The annual habit is found in 196 (24.7%) native genera and 53 (31.2%) native plant families (Table 1). The frequency of the annual habit is significantly higher among dicotyledons than among monocotyledons at the species, generic and familial levels (Table 1). The ten largest genera for annuals in continental Chile are: *Calandrinia* (Portulacaceae) - 31 spp.; *Cryptantha* (Boraginaceae) - 29 spp; *Cristaria* (Malvaceae) - 27 spp.; *Viola* (Violaceae) - 26 spp; *Loasa* (Loasaceae)- 20 spp.; *Chaetanthera*

Stipa

annua Mez

Trichoneura

weberbaueri Pilger

Vulpia

antofagastensis Parodi

australis (Nees) Blom

erolepis (Desv.) Blom

TABLE 1. Summary of frequency of annual habit in the native flora of continental Chile

	Nº species studied				Total N	species %	Genera		Families	
		Obligate annuals	Facultat. annuals	Probably annuals			N	%	N	%
ANGIOSPERMAE	4282	592	79	20	691	16.14	195	26.17	52	34.90
Dicotyledoneae	3374	554	79	19	652	19.32	171	29.38	47	37.90
Monocotyledoneae	908	38	0	1	39	4.30	24	14.72	5	20.00
GYMNOispermae	16	0	0	0	0	0	0	0	0	0
PTERIDOPHYTA	110	1	0	0	1	0.91	1	2.56	1	5.88
TOTAL VASCULAR FLORA	4408	593	79	20	692	15.70	196	24.72	53	31.18

G = 150.34*** (dicots. versus monocots. - species level); G = 15.42*** (dicots. versus monocots. - generic level); G = 3.090; NS (dicots. versus monocots. - familial level).

(Asteraceae) - 18 spp.; *Lepidium* (Cruciferae) - 15 spp.; *Plagiobothrys* (Boraginaceae) and *Vicia* (Papilionaceae) - 14 spp.; *Oenothera* (Onagraceae) - 13 spp. However it should be noted that many of the annuals cited for *Vicia* require further confirmation. Moreover *Calandrinia*, *Cristaria*, *Viola*, *Loasa*, *Plagiobothrys* and *Vicia* require revision. The final number of annual species in these genera judging by the trend seen in other recently revised Chilean genera, might turn out to be lower than presently indicated. Other important genera for annuals in Chile are *Schizanthus* (Solanaceae) with 12 species, and *Gnaphalium* (Asteraceae), *Schizopetalon* (Cruciferae), *Oxalis* (Oxalidaceae), and *Adesmia* (Papilionaceae), all with 10 species.

358 (51.7%) annual species are fairly certainly endemic to continental Chile, while an additional 18 species (2.6%) are characterized by endemic Chilean infraspecific taxa. A further 20 species are probably endemic to continental Chile as are varieties of 3 species. Recent work (Marticorena, unpublished) has shown that 67 (8.5%) of native genera are endemic to continental Chile. Interestingly, seven endemic genera (10.4%) are exclusively annual to facultatively annual in habit (*Cyphocarpus* (Campanulaceae); *Microphyes* (Caryophyllaceae); *Moscharia* (Asteraceae); *Agallis* (Cruciferae); *Scyphanthus* (Loasaceae); *Homalocarpus* (Umbelliferae); *Araeoandra* (Vivianaceae)). These genera, all small, undoubtedly evolved very recently in Chile, as of the Tertiary with the development of arid climates (Arroyo et al., 1988). Such genera constitute interesting material for detailed evolutionary studies in that their ancestors are very likely to be found directly in the extant Chilean flora.

Continental Chile clearly possesses proportionately fewer annuals than the State of California (Raven & Axelrod, 1978) yet more than the Cape floristic region of South Africa (Goldblatt, 1978). Because continental Chile extends further into higher and lower latitudes than California, relatively fewer annuals in continental Chile in relation to California is not entirely unexpected (cf. Arroyo et al., 1988). California is more perfectly matched physiographically and climatically to central

Chile. Work is now in progress in order to determine whether central Chile contains proportionately as many native annuals as California.

ACKNOWLEDGMENTS

Research supported by FONDECYT Grant N° 88-1177. The following persons are thanked for supplying or checking information compiled by us; J. Grau (*Cryptantha*), D. Ford (*Calandrinia*), T. Lammers (Campanulaceae), O. Matthei (Gramineae), T. Bayer (*Alstroemeria*), P. Peñailillo (*Bromus*), C. Von Bohlen (*Mimulus*).

LITERATURE CITED

- ADAMSON, R. S. 1927. The plant communities of Table Mountain. I. Preliminary account. J. Ecol. 15: 275-309.
- ARROYO, MARY T., KALIN, MARTICORENA C. & VILLAGRAN, C. 1984. La flora de la Cordillera de los Andes en el área de Laguna Grande y Laguna Chica, III Región. Chile. Gayana (Botánica) 41: 3-46.
- ARROYO, MARY T., KALIN, SQUEO, F., ARRESTO, J. & VILLAGRAN C. 1988. Effects of aridity on plant diversity in the northern Chilean Andes. Ann. Missouri Bot. Garden 75: 55-78.
- ARROYO, MARY T., KALIN, C., MARTICORENA, P., MIRANDA, O., MATTHEI, A., LANDERO and SQUEO, F., 1989. Contribution to the high elevation flora of the Chilean Patagonia: a checklist of species on mountains on an east-west transect at latitude 50°S. Gayana (Botánica) 46 (1-2): 121-151.
- ARROYO, MARY T., KALIN & SQUEO, F., 1990. Relationship between plant breeding systems and pollination. In (ed.), S. Kawano, Biological Approaches and Evolutionary Trends in Plants. Academic Press, London. pp. 205-227.
- ARROYO, MARY T., KALIN & USLAR, P., 1990. Breeding systems in a temperate mediterranean-type climate montane sclerophyllous forest in central Chile. Bot. J. Linn. Soc. (London): submitted.
- CORREA, M. N. (ed.), 1971. Flora Patagónica. Parte VII. Compositae. Colección Científica del INTA, Buenos Aires.
- DANIN, A. & ORSHAN, G., 1990. The distribution of Raunkiær life forms in Israel in relation to environment. J. Veget. Sci. 1: 41-48.
- DEMPSTER, L. T., 1981. The genus *Galium* (Rubiaceae) in South America II. Allertonia 4: 393-426.
- DI CASTRI, F. & HAJEK, R., 1976. Bioclimatología de Chile. Universidad Católica de Chile, Santiago.
- DI CASTRI, F. 1988. Mediterranean-type shrublands of the world. In (eds.), F. Di Castri, D.W. Goodall & R.L.

- Specht, Mediterranean-type Shrublands, Elsevier, Amsterdam.
- EIG, A. 1931. Les éléments et les groupes phytogeographiques auxiliaires dans la flore palestinienne. Part. I. Feddes Repert. Specierum Nov. Regni Veg. Beih. 63 (1): 1-201.
- EIG, A. 1932. Les éléments et les groupes phytogéographiques auxiliaires dans la flore palestinienne. Parte II. Feddes Repert. Specierum Nov. Regni Veg. Beih. 63 (2): 1-120.
- GOLDBLATT, P. 1978. An analysis of the flora of southern Africa: its characteristics, relationships and origins. Ann. Missouri. Bot. Gard. 65: 369-436.
- MARTICORENA, C. & QUEZADA, M. 1985. Catálogo de la flora vascular de Chile. Gayana (Botánica) 42: 5-157.
- MOONEY, H. A., KUMMEROW, J., JOHNSON, A.W., PARSONS, A. W., KEEFAY, S., HOFFMANN, A., HAYES, R. L., GILBERTO, J., & CHU, C., 1977. The producers - their resources and adaptative responses. In (ed.), H.
- Mooney, Convergent Evolution in Chile and California. Dowden, Hutchinson & Ross, Inc. Stroudsburg, pp. 85-143.
- MOORE, D. M. 1982. Flora of Tierra del Fuego. Anthony Nelson, England.
- RAVEN, PETER H. & AXELROD, D. I. 1978. Origins and relationships of the California flora. Univ. Calif. Publ. Bot. 72: 1-134.
- REICHE, C. 1896-1911. Flora Crítica de Chile. Imprenta Cervantes, Santiago, Chile. 6 vols.
- SCHLEGEI, F., 1966. Pflanzensoziologische und floristische Untersuchungen über Hartlaubgehölze im La Plata-Tal bei Santiago de Chile. Bericht. der Oberhess. Gesell. Natur. Heilk. Giessen, Naturw, Abteil. 34: 183-204.
- SHREVE, F. & WIGGINS, I. M. 1964. Vegetation and Flora of the Sonoran Desert. Stanford Univ. Press, Stanford, California.

Fecha de publicación: 30 abril 1991